



News Articles Videos  
Health & Medicine Mind & Brain Plants & Animals



## Science News

Share Blog Cite

Print Email Bookmark

### All Types Of Antipsychotic Drugs Increase The Risk Of Stroke, Study Finds

*ScienceDaily* (Sep. 2, 2008) — All drugs used to treat psychosis are linked to an increased risk of stroke, and dementia sufferers are at double the risk, according to a study published on the British Medical Journal website.

See also:

#### Health & Medicine

- Elder Care
- Stroke Prevention
- Heart Disease

#### Mind & Brain

- Stroke
- Caregiving
- Schizophrenia

#### Reference

- Multi-infarct dementia
- Dopamine hypothesis of schizophrenia
- Psychopharmacology
- COX-2 inhibitor

Previous research has shown that second generation (atypical) antipsychotic drugs can increase the chances of patients having a stroke. But the risk of stroke associated with first generation (typical) antipsychotics, and whether the risk differs in people with and without dementia, is not known.

Concerns about an increased risk of stroke among people taking atypical antipsychotic drugs were first raised in 2002, particularly in people with dementia. In 2004, the UK's Committee on Safety of Medicines recommended that these drugs should not be used in people with dementia, despite a lack of clear evidence.

A team of researchers from the London School of Hygiene and Tropical Medicine, examined data from the General Practice Research Database (GPRD), which contains the clinical information of more than six million patients registered at over 400 general practices in the UK.

They assessed the effect of exposure to antipsychotic medication on the incidence of stroke in 6 790 patients with a recorded incident of stroke and at least one prescription of any antipsychotic between January 1988 and the end of 2002.

The authors found that during periods when patients were receiving an antipsychotic drug they were 1.7 times more likely to have a stroke, whereas people with dementia were 3.5 times more likely to have a stroke whilst taking any antipsychotic.

The likelihood of having a stroke was slightly higher for people taking atypical antipsychotics than people taking typical antipsychotics.

The study did not look at the specific mechanisms linking antipsychotics and stroke or why the risk is greater with atypical antipsychotics.

Previously, the risk of stroke associated with typical antipsychotics was unclear, say the researchers, but "we have established that all types of antipsychotics carry an increased risk, although the risk might be somewhat higher with the atypical drugs."

They conclude: "We reaffirm that the risks associated with antipsychotic use in patients with dementia generally outweigh the potential benefits, and in this patient group, use of antipsychotic drugs should be avoided wherever possible."

Adapted from materials provided by *BMJ-British Medical Journal*, via *EurekAlert!*, a service of AAAS.

Need to cite this story in your essay, paper, or report? Use one of the following formats:

- APA BMJ-British Medical Journal (2008, September 2). All Types Of Antipsychotic Drugs Increase The Risk Of Stroke, Study Finds. *ScienceDaily*. Retrieved September 3, 2008, from <http://www.sciencedaily.com/releases/2008/08/080828220511.htm>
- MLA

### Search ScienceDaily

Number of stories in archives: 44,032

Find with keyword(s):  Search

Enter a keyword or phrase to search ScienceDaily's archives for related news topics, the latest news stories, reference articles, science videos, images, and books.

Ads by Google

Advertise here

**The Theory of Everything**  
Read new perspectives on quantum physics, relativity, and cosmology.

**Treat Alzheimer's Disease**  
Time tested remedies. Most effective since Safe effective.

**Why Your Belly is Fat**  
I fought with excess belly fat for years until I found this diet.

Just In:

Genetic Link To Relationship Difficulties Found

### Science Video News



#### Stopping Strokes

To prevent strokes in at-risk patients who suffer from atrial fibrillation, an abnormal heart rhythm, researchers have devised a mechanical. ... > [full story](#)

[Pediatric Neurologists Use MRI to Understand How Strokes Impair Verbal Abilities](#)

[Psychiatrists Can Predict Onset of Alzheimer's with New EEG Test](#)

[Interventional Neuroradiologists Treat Brain Strokes with New Kind of Stent](#)

[more science videos](#)



### Breaking News

... from NewsDaily.com

[Bush visits Louisiana as state claws back from storm](#)

[Virus is passed from parent to child in the DNA](#)

[Clones' offspring may be in food supply: FDA](#)

[Marriage problems? Husband's genes may be to blame](#)

[EU lawmakers open battle against sheep, goat tags](#)

[more science news](#)

### In Other News ...

[North Korea reassembling nuclear facility: report](#)

[Afghans accept U.S. probe offer on civilian deaths](#)

[Suspected U.S. troops kill 20 in Pakistan attack](#)

[Shooting spree leaves six dead in Washington state](#)

[Storms swirl in Atlantic, deadly floods hit Haiti](#)

[Massive Canada Arctic ice shelf breaks away](#)

[U.S. urges probe of journalist's death in Ingushetia](#)

[Pakistani indicted in NY for attack on U.S. soldiers](#)  
[more top news](#)



Yes. You do not have to accept nausea or vomiting caused by chemotherapy...



Roll over for the full answer.

Copyright Reuters 2008. See [Restrictions](#).

hourly updated newsfeeds in your RSS reader:

[Email Newsletters](#)

[RSS Newsfeeds](#)

**Feedback** *... we want to hear from you!*

Tell us what you think of the new ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

Your Name:

Your Email:

Comments:

Click button to submit feedback:

[About This Site](#) | [Editorial Staff](#) | [Awards & Reviews](#) | [Contribute News](#) | [Advertise With Us](#) | [Privacy Policy](#) | [Terms of Use](#)  
Copyright © 1995-2008 ScienceDaily LLC — All rights reserved — Contact: [editor@sciencedaily.com](mailto:editor@sciencedaily.com)